

# Coherent states of non-Hermitian quantum systems

B. Roy, P. Roy\*

*Physics & Applied Mathematics Unit, Indian Statistical Institute, Kolkata 700 108, India*

Received 23 March 2006; received in revised form 11 May 2006; accepted 5 June 2006

Available online 13 June 2006

Communicated by P.R. Holland

---

## Abstract

We use the Gazeau–Klauder formalism to construct coherent states of non-Hermitian quantum systems. In particular we use this formalism to construct coherent state of a  $\mathcal{PT}$  symmetric system. We also describe the construction of coherent states following Klauder’s minimal prescription.

---

## 1. Introduction

In recent years non-Hermitian quantum mechanics have been extensively studied from various stand points [1]. Among the different non-Hermitian systems, there is a class of problems which are  $\mathcal{PT}$  invariant,  $\mathcal{P}$  and  $\mathcal{T}$  being the parity and time reversal operator respectively and it was shown that some of the non-Hermitian  $\mathcal{PT}$  symmetric problems admit real eigenvalues [2]. Subsequently it was pointed out that the reality of the spectrum is essentially due to  $\eta$ -pseudo Hermiticity [3]. Interestingly some of the  $\eta$ -pseudo-Hermitian systems are also  $\mathcal{PT}$  symmetric. Because of this property and also because of their intrinsic interest  $\mathcal{PT}$  symmetric and  $\eta$ -pseudo-Hermitian potentials have been examined widely [4,5].

On the other hand coherent states play an important role in the context of Hermitian quantum mechanics [6]. Recently the concept of coherent states was also introduced to  $\mathcal{PT}$  symmetric quantum mechanics [7]. However, as in Hermitian models, coherent states corresponding to arbitrary non-Hermitian potentials are not easy to construct. This is mainly due to the fact that symmetry of the problem i.e., a knowledge of the ladder operators may not always be known. This difficulty can however be overcome by using the Gazeau–Klauder (GK) approach [8]. Recently Klauder [9] has suggested a set of requirements which a coherent state should satisfy and he proposed a method of con-

structing coherent states for solvable potentials. This method is simple and has been applied successfully to a number of exactly solvable Hermitian potentials with discrete [10] as well as continuous spectrum [8]. Here our objective is to show that with an appropriately chosen inner product the GK formalism can also be extended to non-Hermitian potentials. In particular we shall use this technique to construct coherent states of the  $\mathcal{PT}$  symmetric Scarf I potential. We shall also construct coherent states of the same potential satisfying a minimal set of requirements.

## 2. Some results on $\mathcal{PT}$ symmetric systems

Let us consider a Hamiltonian  $H$  such that

$$H\psi_n(x) = E_n\psi_n(x) = \omega e_n\psi_n, \quad e_0 = 0. \quad (1)$$

Then  $H$  is said to be  $\mathcal{PT}$  symmetric if

$$H = \mathcal{PT}H\mathcal{PT}, \quad (2)$$

where

$$\left. \begin{aligned} \mathcal{P}: & \quad x \rightarrow -x, \quad p \rightarrow -p \\ \mathcal{T}: & \quad x \rightarrow x, \quad p \rightarrow -p, \quad i \rightarrow -i \end{aligned} \right\}. \quad (3)$$

Furthermore, if in addition to (2) the wave functions are also  $\mathcal{PT}$  invariant i.e.,

$$\mathcal{PT}\psi_n = \pm\psi_n, \quad (4)$$

then  $\mathcal{PT}$  symmetry is said to be unbroken. On the other hand if the Hamiltonian is  $\mathcal{PT}$  invariant while the wave functions

---

\* Corresponding author.

E-mail addresses: barnana@isical.ac.in (B. Roy), pinaki@isical.ac.in (P. Roy).

are not, then  $\mathcal{PT}$  symmetry is said to be spontaneously broken. It may be noted that systems with spontaneously broken  $\mathcal{PT}$  symmetry are characterised by complex conjugate pair of eigenvalues. Here we shall only consider systems with unbroken  $\mathcal{PT}$  symmetry.

A major difference between the non-Hermitian theories and the Hermitian ones lies in the definition of inner product. In the case of  $\mathcal{PT}$  symmetric systems neither the standard definition of inner product nor the straightforward generalisation, namely,

$$\langle \psi_m | \psi_n \rangle^{\mathcal{PT}} = \int [\mathcal{PT} \psi_m] \psi_n dx = (-1)^n \delta_{mn} \quad (5)$$

work because the norm becomes negative for some of the states. Consequently a modification is necessary so that the norm is always positive. Indeed it has been shown that for  $\eta$ -pseudo-Hermitian quantum mechanics it is possible to define an inner product which is positive definite i.e., the Hamiltonian is Hermitian with respect to that inner product [3]. For  $\mathcal{PT}$  symmetric cases a convenient option is to use the  $\mathcal{CPT}$  norm. For  $\mathcal{PT}$  symmetric theories with unbroken  $\mathcal{PT}$  symmetry the  $\mathcal{CPT}$  inner product is defined by [11]

$$\langle \psi_m | \psi_n \rangle^{\mathcal{CPT}} = \int [\mathcal{CPT} \psi_m] \psi_n dx = \delta_{mn}, \quad (6)$$

where  $\mathcal{C}$  is called the charge operator and is defined by [11]

$$\mathcal{C}(x, y) = \sum_{n=0}^{\infty} \psi_n(x) \psi_n(y). \quad (7)$$

The action of the charge operator on the eigenfunctions is given by

$$\mathcal{C} \psi_n(x) = \int \mathcal{C}(x, y) \psi_n(y) dy = (-1)^n \psi_n(x). \quad (8)$$

Another property which would be very useful later is the completeness of eigenfunctions. In coordinate space the completeness property can be expressed in terms of the charge operator as

$$\sum_{n=0}^{\infty} \psi_n(x) [\mathcal{CPT} \psi_n(y)] = \sum_{n=0}^{\infty} (-1)^n \psi_n(x) \psi_n(y) = \delta(x - y). \quad (9)$$

### 3. GK formalism for $\mathcal{PT}$ symmetric systems

It is known that coherent states can be constructed using different techniques (e.g., a coherent state may be defined as a minimum uncertainty state, annihilation operator eigenstate, etc.) and usually they have different properties. In the GK formalism a coherent state should satisfy the following criteria [8]:

- (1) continuity of labelling;
- (2) temporal stability;
- (3) resolution of identity;
- (4) action identity.

For  $\mathcal{PT}$  symmetric systems a GK coherent state is a two parameter state defined by [8,9]

$$\psi(x; J, \gamma) = \frac{1}{\sqrt{\mathcal{N}(J)}} \sum_{n=0}^{\infty} \frac{J^{\frac{n}{2}} \exp(-i\gamma e_n)}{\sqrt{\rho_n}} \psi_n(x), \quad (10)$$

where  $J \geq 0$  and  $-\infty < \gamma < +\infty$  are two parameters and  $\psi_n(x)$  are the eigenstates. In (10),  $\rho_n$  are a set of numbers defined as

$$\rho_0 = 1, \quad \rho_n = \prod_{i=1}^n e_i. \quad (11)$$

It is now necessary to determine the normalisation constant  $\mathcal{N}(J)$ . Now using the  $\mathcal{CPT}$  inner product (6) the normalisation constant  $\mathcal{N}(J)$  can be obtained from the condition  $\langle \psi(x; J, \gamma) | \psi(x; J, \gamma) \rangle^{\mathcal{CPT}} = 1$ :

$$\mathcal{N}(J) = \sum_{n=0}^{\infty} \frac{J^n}{\rho_n}, \quad 0 < J < R = \lim_{n \rightarrow \infty} (\rho_n)^{\frac{1}{n}}. \quad (12)$$

Let us now examine whether or not the coherent states (10) satisfy the criteria mentioned above. From the construction it is clear that the coherent states are continuous in their labels i.e.,  $(J, \gamma) \rightarrow (J', \gamma') \Rightarrow \psi(x; J, \gamma, x) \rightarrow \psi(x; J', \gamma')$ . It may also be noted that although  $H$  is not Hermitian, the time evolution operator is still given by  $e^{-iHt}$ . Using (1) and (10) the pseudo-unitary time evolution is found to be

$$e^{-iHt} \psi(x; J, \gamma) = \psi(x; J, \gamma + \omega t). \quad (13)$$

From (13) we find that the GK coherent states are temporally stable. We now show that the coherent state also admits a resolution of unity:

$$\begin{aligned} & \int d\mu(\gamma, J) [\mathcal{CPT} \psi(y; J, \gamma)] f(J) \psi(x; J, \gamma) \\ &= \lim_{\Gamma \rightarrow \infty} \frac{1}{2\Gamma} \int_{-\Gamma}^{\Gamma} d\gamma \int_0^{\infty} [\mathcal{CPT} \psi(y; J, \gamma)] \psi(x; J, \gamma) f(J) dJ \\ &= \sum_{n=0}^{\infty} \frac{(-1)^n \psi_n(y) \psi_n(x)}{\rho_n} \int J^n \frac{f(J)}{\mathcal{N}(J)} dJ = \delta(x - y) \end{aligned} \quad (14)$$

if the following moment problem has a solution:

$$\rho_n = \int_0^{\infty} J^n \frac{f(J)}{\mathcal{N}(J)} dJ. \quad (15)$$

Thus we find that the GK coherent state for  $\mathcal{PT}$  symmetric systems provide a resolution of unity subject to the solution of the moment problem (15).

We now proceed to check the action identity. Using (12) it can be easily checked that

$$\begin{aligned} & \langle \psi | H | \psi \rangle^{\mathcal{CPT}} \\ &= \frac{1}{\mathcal{N}(J)} \sum_{m, n=0}^{\infty} \frac{J^{(n+m)/2} e^{-i\gamma(e_n - e_m)} \omega e_n}{\sqrt{\rho_m \rho_n}} (-1)^n \langle \psi_m | \psi_n \rangle \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{\mathcal{N}(J)} \sum_{m,n=0}^{\infty} \frac{J^{(n+m)/2} e^{-i\gamma(e_n - e_m)} \omega e_n}{\sqrt{\rho_m \rho_n}} (-1)^{n+m} \delta_{mn} \\
 &= \omega J \tag{16}
 \end{aligned}$$

so that the criteria (4) is also satisfied. Thus the coherent state (10) satisfy all the criteria (1)–(4). It may be noted the construction ultimately boils down to a solution of the moment problem (15).

**4. GK coherent states for  $\mathcal{PT}$  symmetric Scarf I potential**

As an example we shall now construct GK coherent state of an exactly solvable  $\mathcal{PT}$  symmetric potential. Thus we consider the  $\mathcal{PT}$  symmetric Scarf I potential [12–14]. The Schrödinger equation is given by

$$\left[ -\frac{d^2}{dx^2} + V(x) \right] \psi_n = E_n \psi_n, \tag{17}$$

where the potential  $V(x)$  is given by

$$\begin{aligned}
 V(x) = & \frac{2(\alpha^2 + \beta^2) - 1}{4} \frac{1}{\cos^2(x)} + \frac{(\alpha^2 - \beta^2)}{2} \frac{\sin(x)}{\cos^2(x)} \\
 & - \frac{(\alpha + \beta + 1)^2}{4}, \quad x \in \left[ -\frac{\pi}{2}, \frac{\pi}{2} \right], \tag{18}
 \end{aligned}$$

where  $\alpha$  and  $\beta$  are complex parameters such that  $\beta^* = \alpha$  and  $\alpha_R > \frac{1}{2}$ . The eigenvalues and the corresponding eigenfunctions are given by [12–14]

$$E_n = \omega e_n = n(n + 2\alpha_R + 1), \quad n = 0, 1, 2, \dots, \tag{19}$$

$$\psi_n(x) = N_n (1 - \sin x)^{\frac{\alpha}{2} + \frac{1}{4}} (1 + \sin x)^{\frac{\beta}{2} + \frac{1}{4}} P_n^{(\alpha, \beta)}(\sin x), \tag{20}$$

where  $N_n$  denotes the normalisation constant and  $P_n^{(\alpha, \beta)}$  are Jacobi polynomials [15]. It may be noted that the Hamiltonian is  $\mathcal{PT}$  symmetric as well as  $\mathcal{P}$ -pseudo-Hermitian:

$$\begin{aligned}
 H &= \mathcal{PT} H \mathcal{PT}, \\
 H^\dagger &= \mathcal{P}^{-1} H \mathcal{P}. \tag{21}
 \end{aligned}$$

Also the wave functions are  $\mathcal{PT}$  invariant:

$$\mathcal{PT} \psi_n(x) = \psi_n(x) \tag{22}$$

and they satisfy the relation

$$\int [CPT \psi_m(x)] \psi_n(x) dx = \delta_{mn}. \tag{23}$$

In this case

$$\begin{aligned}
 e_n &= n(n + v), \quad \rho_0 = 1, \\
 \rho_n &= \frac{\Gamma(n + 1)\Gamma(n + v + 1)}{\Gamma(v + 1)}, \quad R = \infty, \tag{24}
 \end{aligned}$$

where  $v = 2\alpha_R + 1$ .

Thus the GK coherent state for the Scarf I potential (in the coordinate representation) is given by

$$\psi(x; J, \gamma) = \frac{1}{\sqrt{\mathcal{N}(J)}} \sum_{n=0}^{\infty} \frac{J^{\frac{\alpha}{2}} \exp(-i\gamma e_n)}{\sqrt{\rho_n}} \psi_n(x), \tag{25}$$

where  $\psi_n(x)$  and  $\rho_n$  are given by (20) and (24), respectively. From (12) the normalisation constant is found to be

$$\begin{aligned}
 \mathcal{N}(J) &= \left[ \Gamma(v + 1) \sum_{n=0}^{\infty} \frac{J^n}{\Gamma(n + 1)\Gamma(n + v + 1)} \right] \\
 &= J^{-v/2} \Gamma(v + 1) I_v(2\sqrt{J}), \tag{26}
 \end{aligned}$$

where  $I_v(x)$  stands for the Bessel function of the first kind [15].

It is now necessary to show that the moment problem (15) has a solution. Using the relation [15]

$$\begin{aligned}
 \int_0^\infty x^\mu K_\delta(ax) dx &= 2^{\mu-1} a^{-\mu-1} \Gamma\left(\frac{1 + \mu + \delta}{2}\right) \Gamma\left(\frac{1 + \mu - \delta}{2}\right), \\
 \text{Re}(\mu + 1 \pm \delta) &> 0, \quad \text{Re}(a) > 0 \tag{27}
 \end{aligned}$$

we find that

$$f(J) = \frac{1}{\pi} I_v(2\sqrt{J}) K_v(2\sqrt{J}) > 0 \tag{28}$$

provides a solution to the moment problem (15). Now it can be shown that

$$\begin{aligned}
 &\int d\mu(J, \gamma) [CPT \psi(y; J, \gamma)] f(J) \psi(x; J, \gamma) \\
 &= \frac{1}{2\pi} \int_0^{2\pi} d\gamma \int_0^\infty dJ \mathcal{N}(J)^{-1} \psi(x; J, \gamma) [CPT \psi(y; J, \gamma)] \\
 &= \sum_{n=0}^{\infty} (-1)^n \psi_n(x) \psi_n(y) = \delta(x - y). \tag{29}
 \end{aligned}$$

Thus the coherent state provide a resolution of unity. It can now be easily shown that the states (25) have many other properties characteristic of coherent states e.g. they are non-orthogonal:

$$\begin{aligned}
 &\langle \psi(x; J', \gamma') | \psi(x; J, \gamma) \rangle^{CPT} \\
 &= \frac{\Gamma(v + 1)}{\sqrt{\mathcal{N}(J)\mathcal{N}(J')}} \sum_{n=0}^{\infty} \frac{(JJ')^{n/2}}{\Gamma(n + 1)\Gamma(n + v + 1)} e^{jn(n+v)(\gamma - \gamma')}. \tag{30}
 \end{aligned}$$

From the above example it is thus clear that subject to the solution of the moment problem (15), it is possible to construct GK coherent states of any exactly solvable  $\mathcal{PT}$  symmetric or  $\eta$ -pseudo-Hermitian potential with real spectrum.

**4.1. Minimal coherent state for non-Hermitian potentials**

In the last section we considered coherent states satisfying four conditions. However, if we relax conditions (2) and (4) and construct coherent states following Klauder’s minimal prescription [16], then a wider class of such states can be generated [17]. In this case a coherent state is required to satisfy two of the four criteria, namely, (1) continuity in labelling and, (2) resolution of identity, mentioned earlier. Thus a coherent state corresponding to (18) is defined as

$$\psi(x; \beta) = \frac{1}{\sqrt{\mathcal{N}(|\beta|)}} \sum_{n=0}^{\infty} \frac{\beta^n}{\sqrt{\rho_n}} \psi_n(x), \tag{31}$$

where  $\beta = re^{i\theta}$  is a complex number and  $\psi_n(x)$  are given by (20). Here  $\rho_n$  are a set of positive constants which would be specified later. The normalisation constant is determined from the condition  $\langle \psi(x; \beta) | \psi(x; \beta) \rangle^{\mathcal{CPT}} = 1$  and is given by

$$\mathcal{N}(|\beta\rangle) = \sum_{n=0}^{\infty} \frac{(\beta\bar{\beta})^n}{\rho_n}. \tag{32}$$

Clearly  $\beta \rightarrow \beta' \Rightarrow \psi(x; \beta) \rightarrow \psi(x; \beta')$  i.e., the coherent state (31) is continuous in the label. Furthermore

$$\int d^2\beta [\mathcal{CPT} \psi(y; \beta)] f(|\beta\rangle) \psi(x; \beta) = 2\pi \sum_{n=0}^{\infty} \frac{(-1)^n \psi_n(y) \psi_n(x)}{\rho_n} \int_0^{\infty} r^{2n+1} \frac{f(r)}{\mathcal{N}(r)} dr. \tag{33}$$

Thus (31) admits resolution of unity if the positive constants  $\rho_n$  are such that the Stieltjes moment problem (15) has a solution i.e.,

$$2\pi \int_0^{\infty} r^{2n+1} \frac{f(r)}{\mathcal{N}(r)} dr = \rho_n. \tag{34}$$

In a recent work [17] solutions to such moment problems for a number of different forms of  $\rho_n$  have been found by using Mellin transform technique. Here we consider a simple example and take  $\rho(n) = \Gamma(2n + 1)$ . In this case we find from (32)

$$\mathcal{N}(r) = \cosh(r). \tag{35}$$

Now using the result

$$\int_0^{\infty} x^{2n} e^{-x} dx = \Gamma(2n + 1) \tag{36}$$

$f(r)$  is found to be

$$f(r) = \frac{1}{2\pi} \frac{e^{-r}}{r}. \tag{37}$$

Since  $f(r) > 0$ , the coherent state

$$\psi(x; \beta) = \frac{1}{\sqrt{\cosh(r)}} \sum_{n=0}^{\infty} \frac{\beta^n}{\sqrt{\Gamma(2n + 1)}} \psi_n(x) \tag{38}$$

with  $\psi_n(x)$  are given by (20), admits a resolution of identity. Also it can be shown that overlap between two coherent states is

$$\langle \psi(x; \beta') | \psi(x; \beta) \rangle^{\mathcal{CPT}} = \frac{\cosh(\sqrt{\beta'\bar{\beta}})}{\sqrt{\cosh(r') \cosh(r)}}. \tag{39}$$

Thus (38) furnishes a new coherent state of the  $\mathcal{PT}$  symmetric Scarf I potential. We note that in contrast to the GK coherent state (25), the minimal coherent state (38) is not temporally stable.

### 5. Conclusion

Here it has been shown that by suitably modifying the normalisation procedure, it is possible to extend the GK formalism to non-Hermitian systems. Furthermore new families of coherent states may also be constructed following Klauder’s minimal prescription. One such state with  $\rho_n = \Gamma(2n + 1)$  has been constructed here. Finally we note that the construction of coherent states outlined here may also be extended to  $\eta$ -pseudo-Hermitian systems which are not necessarily  $\mathcal{PT}$  symmetric [18].

### References

- [1] N. Moiseyev, M. Glück, Phys. Rev. E 63 (2001) 041103; D.R. Nelson, M. Shnerb, Phys. Rev. E 58 (1998) 1383; N. Hatano, D.R. Nelson, Phys. Rev. B 38 (1998) 8384; E. Narevicius, P. Serra, N. Moiseyev, Europhys. Lett. 62 (2003) 789.
- [2] C.M. Bender, S. Boettcher, Phys. Rev. Lett. 80 (1998) 5243.
- [3] A. Mostafazadeh, J. Math. Phys. 43 (2002) 205; A. Mostafazadeh, J. Math. Phys. 43 (2002) 2824; A. Mostafazadeh, J. Math. Phys. 43 (2002) 3944; A. Mostafazadeh, J. Math. Phys. 44 (2003) 974.
- [4] C.M. Bender, S. Boettcher, P.N. Meisinger, J. Math. Phys. 40 (1999) 2201; M. Znojil, J. Phys. A 32 (1999) 4653; G. Levai, M. Znojil, J. Phys. A 33 (2000) 7165; B. Bagchi, C. Quesne, Phys. Lett. A 273 (2000) 285; A. Mostafazadeh, J. Math. Phys. 43 (2002) 205; A. Mostafazadeh, J. Math. Phys. 43 (2002) 2814; A.A. Andrianov, F. Cannata, J.P. Dedonder, M.V. Ioffe, Int. J. Mod. Phys. A 14 (1999) 2675.
- [5] A. Mostafazadeh, J. Math. Phys. 46 (2005) 102108; A. Mostafazadeh, J. Phys. A 38 (2005) 3213; A. Mostafazadeh, J. Phys. A 38 (2004) 11645.
- [6] A.M. Perelemov, Generalised Coherent States and Some of their Applications, Springer-Verlag, Berlin, 1986; J.R. Klauder, B.-S. Skagerstam, Coherent States: Applications in Physics and Mathematical Physics, World Scientific, Singapore, 1985.
- [7] B. Bagchi, C. Quesne, Mod. Phys. Lett. A 16 (2001) 2449; See also, M. Znojil, hep-th/0012002.
- [8] J.-P. Gazeau, J.R. Klauder, J. Phys. A 32 (1999) 123.
- [9] J.R. Klauder, J. Phys. A 29 (1996) L293.
- [10] J.-P. Antoine, J.-P. Gazeau, P. Monceau, J.R. Klauder, K.A. Penson, J. Math. Phys. 42 (2001) 2349; M. Novaes, J.-P. Gazeau, J. Phys. A 36 (2003) 199; A. Chenaghloou, H. Fakhri, Mod. Phys. Lett. A 17 (2002) 1701; A.H. El Kinani, M. Daoud, Phys. Lett. A 283 (2001) 291; M. Roknizadeh, M.K. Tavassoly, J. Math. Phys. 46 (2005) 042110.
- [11] C.M. Bender, D.C. Brody, H.F. Jones, Phys. Rev. Lett. 89 (2002) 27041; C.M. Bender, D.C. Brody, H.F. Jones, Phys. Rev. Lett. 92 (2004) 119902.
- [12] G. Levai, J. Phys. A, in press.
- [13] G. Levai, M. Znojil, J. Phys. A 33 (2000) 7165.
- [14] G. Levai, M. Znojil, Mod. Phys. Lett. A 16 (2001) 1973.
- [15] I.S. Gradshteyn, I.M. Ryzhik, in: A. Jeffrey (Ed.), Tables of Integrals, Series and Products, Academic Press, San Diego, 1994.
- [16] J.R. Klauder, J. Math. Phys. 4 (1963) 1035; J.R. Klauder, J. Math. Phys. 4 (1963) 1058.
- [17] J.R. Klauder, K.A. Penson, J.-M. Sixdeniers, Phys. Rev. A 64 (2001) 013817.
- [18] Z. Ahmed, Phys. Lett. A 290 (2001) 19.