

PROFESSOR RONALD AYLMEY FISHER

EARLY DAYS.

Ronald Aylmer Fisher was born on the 17th of February, 1890, in East Finchley, one of the northern suburbs of London, where his father had a large establishment. He was the youngest of seven children which would have been eight if his twin brother had lived. His father, G. Fisher, of the well-known firm of auctioneers, Robinson and Fisher in King Street, St. James's, was a man of extraordinary energy and had a wide and detailed knowledge of the fine arts. The father's family were mostly businessmen; but an uncle, the younger brother of his father, was placed high as a Cambridge Wrangler and entered the Church. The mother's father was a successful London solicitor noted for his social qualities. Besides sturdy common sense and interest in practical affairs, taste for mathematics and the fine arts, Fisher inherited a spirit of adventure from his family. His mother's only brother threw up excellent prospects in London to collect wild animals in Africa, and one of his brothers returned from the Argentine to serve in the European War and was killed in 1915. Extremely defective vision saved Fisher himself from a similar fate, for, though in 1914 he had qualified himself for commission, he was refused throughout the war the opportunity of military service.

As with many mathematicians, Fisher's special ability showed at an early age. Before he was six, his mother was reading him a popular book on Astronomy, an interest which he followed eagerly during boyhood. Love of mathematics dominated his educational career; and he was fortunate in coming under the tuition of a brilliant mathematical teacher, W. N. Roe of Stanmore Park, also well-known in England as a Somersetshire cricketer. At Harrow he worked under C. H. P. Mayo and W. N. Roseveare. The peculiar circumstances of the teaching by the latter will be of interest to statisticians familiar with Fisher's geometrical methods. On account of his eyes he was forbidden to work by artificial light, and when he would go to work with Mr. Roseveare in the evenings the instruction was given purely by ear without the use of paper and pencil or any other visual aid. To this early training may be attributed Fisher's tendency to use hypergeometrical representation, and his great power in penetrating into problems requiring geometrical intuition. It is, perhaps, a consequence of this that throughout life his solutions have been singularly independent of symbolism. He does not usually attempt to write down the analysis until the problem is solved in his mind, and sometimes, he confesses, after the key to the solution has been forgotten.

CAMBRIDGE.

Though born to sufficiently wealthy parents, owing to financial difficulties of the family at the time, it would have been difficult for him to proceed to the University without the aid of scholarships which he succeeded in winning. He joined the Gonville and Caius College, Cambridge, in 1909; and passed the Mathematical Tripos Part II in 1912 as a Wrangler with distinction in the optical papers in schedule B. He spent another year at Cambridge with a Studentship in Physics, and studied statistical mechanics and quantum theory under James Jeans, and the theory of errors under F. J. M. Stratton.

At this period he was consciously out of sympathy with certain tendencies prevalent at the time in the University. Under the influence of Bateson it was widely believed that recent work on genetics had discredited the evolutionary principles developed by Darwin. Fisher became keenly interested in Mendelian evidence, and was convinced that this furnished a basis for a quantitative and mathematical treatment of the theory of evolution. He began at this early date a series of studies of which the outside world knew nothing until the publication in 1930 of his book on "The Genetical Theory of Natural Selection."

The second influence at Cambridge with which Fisher found himself out of sympathy was the recent passage of control in mathematical teaching from the earlier tradition

of mathematical physicists to a school of pure mathematicians of largely continental derivation. The explicit statement of a rigorous argument interested him, but only on the important condition that such explicit demonstration of rigour was needed. Mechanical drill in the technique of rigorous statement was abhorrent to him, partly for its pedantry, and partly as an inhibition to the active use of the mind. He felt it was more important to think actively, even at the expense of occasional errors from which an alert intelligence would soon recover, than to proceed with perfect safety at a snail's pace along well-known paths with the aid of the most perfectly designed mechanical crutches. The real interpretation of his state of mind at this time is probably that he was fitting himself for mathematical research rather than for mathematical teaching. Fisher himself thinks that he was merely a very wilful and impatient youngman. This is no doubt true, but he was impatient not because he was young but because he was a creative genius.

EARLY WORK IN STATISTICS.

Although he had a successful academic career at Cambridge he had not yet found his vocation in life. I have noted before that for some time he studied mathematical physics. He also attended, it is said, one lecture on statistics under G. Udny Yule. He read at this time very carefully and very critically Karl Pearson's "Mathematical Contributions to the Theory of Evolution." Shortly before the War, in 1914, he first came into contact with Karl Pearson. H. E. Soper had discussed in a paper in the *Biometrika* the distribution of the coefficient of correlation in samples from an infinite, normally distributed, bi-variate population, but had succeeded in giving only approximate results by a tedious process. The problem, which was clearly stated by Soper, was one after Fisher's own heart, and in a week he had sent to Karl Pearson the exact solution but in a more hastily and roughly sketched form than even that in which it was subsequently published in the *Biometrika*.¹ This first paper, which was the starting point of the modern theory of exact sampling distributions, shows in a characteristic manner the working of Fisher's genius.

It was only after he had completed this important work that Fisher read the earlier papers of "Student",² and noticed how the representation in many-dimensional space so perfectly supplied the completion of "Student's" tentative ideas. A little later he solved the problem of the exact distribution of the intra-class correlation. This solution together with his z -transformation for coefficients of correlation of both kinds was published in 1921 in *Metron*.³

Fisher was now seriously interested in theoretical statistics, and any invitation to engage in academic work of a statistical character would have been no doubt welcome, but no such opportunity occurred for some considerable time. During the war he was at first engaged in statistical work in the office of the Mercantile and General Investment Company from 1913 to 1915, and from 1915 to 1919 in teaching physics and mathematics in public schools. It cannot be said that school teaching was congenial to him, but the few years spent in such work was far from wasted. Experience in teaching physics supplied what Fisher has always regarded as the most serious omission of his studies at Cambridge, namely, actual touch with problems of practical experimentation. On the whole therefore it was probably fortunate that the opportunity of independent statistical work came at a time when his outlook was more mature.

ROTHAMSTED EXPERIMENTAL STATION.

At the close of the War he was offered the post of the chief statistician under Karl Pearson at the Galton Laboratory. At about the same time Sir John Russell, Director of the Rothamsted Agricultural Station, offered him the post of statistician with opportunities for building up a Statistical Laboratory at Rothamsted. Neither post had much financial attraction, but he unhesitatingly accepted the Rothamsted offer as he thought that facilities for independent research would be greater there.

1. *Biometrika*, x, 507-521 (1915).

2. W. S. Gosset, whose untimely death in November, 1937, took away a great pioneer in statistics.

3. *Metron*, i, part 4, 1-32.

At Rothamsted Fisher found his real vocation for life, and there followed a series of years of intense activity. On the theoretical side he continued his work on sampling distributions and in 1921 completed the classical memoir on "Mathematical Foundations of Theoretical Statistics"¹⁸ which was intended to supply the framework for modern statistical theory. On the applied side the method for separating the slow changes in time series was given in the same year,⁴ and a little later in 1924, the important memoir on the influence of rainfall on the yield of wheat.⁵ In 1923 was published the first paper on field trials⁶ which led to a revolution in the technique of agricultural trials throughout the world, and was the starting point of the work on the design of experiments. The z-test of significance and the arithmetical procedure known as the 'Analysis of Variance' took their present form at about the same time.

"Statistical Methods for Research Workers" was first published in 1925, and has since then run to six editions. This book has probably done more than anything else to make research workers in most diverse fields of study familiar with the practical applications of modern statistical methods, and to create a statistical attitude of mind among the younger generation of scientists.

During the next few years we have a number of important papers on exact sampling distributions, the theory of estimations, the Chi-square measure of contingency, Bayes' theorem and inverse probability on the mathematical side, and the arrangement of field experiments on the applied side.

GENETICAL THEORY OF NATURAL SELECTION.

Fisher had been working on Mendelism and genetics for a long time and occasional papers had been published since 1918, but from about 1925 this subject began to engage increasing attention. The book on "Genetical Theory of Natural Selection" was published in 1930, and constituted a landmark in the history of the subject. A mathematical theory was developed on the basis of recent genetical researches to establish the principle of Natural Selection, on a more rigorous basis than Darwin had claimed, as the effective cause of evolutionary change. This principle is considered, not merely as a qualitative hypothesis in connexion with the classes of facts which it is capable of explaining, but rather in the manner of theoretical physics, as an agent which is bound to cause definite changes at a calculable rate. Factors recognized by Darwin, such as sexual selection, can then be examined on a quantitative rather than a speculative basis; and phenomena unknown to Darwin, such as genetical 'dominance' and the selection of fertility in human societies are found to illustrate unexpected effects of the evolutionary mechanism. Like Clerk-Maxwell who translated Faraday's concepts into mathematical language and developed the electromagnetic theory of light, Fisher gave a quantitative form to Darwin's views and built up a statistical theory of evolution.

ACADEMIC HONOURS.

The course of events in his life was in the meantime following a normal channel. He was married in 1917 to Ruth Eileen, daughter of H. Grattan Guinness, M.D., and several children were born to them. In 1920 he was elected a Fellow of Gonville and Caius College, Cambridge, and was awarded the Sc.D. degree of the same University in 1926. The influence of his work was in the meantime spreading rapidly, and he was elected a Fellow of the Royal Society in 1929, and was awarded the Weldon Medal in the same year. He was elected Honorary Fellow of the American Statistical Association in 1930, and Foreign Member of the American Academy of Arts and Science in 1934.

GALTON PROFESSOR IN LONDON.

Fisher was appointed Galton Professor in the University of London on the retirement of Karl Pearson in 1933. He has always been interested in eugenics, and has been

4. *Jour. Agri. Sc.* xi, 107-135.

5. *Phil. Trans. Roy. Soc. B.* cccxiii, 89-142.

6. *Jour. Agri. Sc.* xiii, 311-320.

for a long time associated with the Eugenics Society first as Honorary Secretary and later as Vice-President. Besides the importance of his work in genetics it was therefore peculiarly appropriate that he should succeed to the Chair created by an endowment from the great founder of the science of eugenics. He also took over from Karl Pearson the editorial charge of the *Annals of Eugenics* from 1933. During the last few years Fisher has devoted a great deal of attention to the application of statistical theory to genetic research and a large number of papers have been published in this subject.

The methods first developed in connexion with agricultural experiments have found in recent years increasing application in industry, medicine and public health, education and psychology, and scientific experiments of all kinds. In 1935 Fisher published a book on "The Design of Experiments" in which the theoretical principles were developed with a great variety of illustrative examples.

In 1936 Fisher visited the United States and received an honorary degree from the Harvard University on the occasion of its Tercentenary celebrations. Early in 1937 he accepted the Honorary Fellowship of the Indian Statistical Institute, and the invitation to preside over the first session of the Indian Statistical Conference in January, 1938.

MAIN CURRENTS OF FISHER'S WORK.

Fisher's work falls naturally into three main streams:—(a) contributions to the mathematical theory of statistics; (b) application of statistical theory to agriculture and the design of experiments; and (c) contributions to genetics.

On the mathematical side, for the first time, Fisher has supplied a unified, and general theory for drawing rigorous conclusions from statistical data. The logical and philosophical consequences of Fisher's theory still require to be fully worked out. Its importance in the theory of knowledge can be easily appreciated from the claim that it supplies the only rigorous logical foundation for all inductive inferences in science.

The theory of design of experiments is intended to supply an adequate technique for collecting the primary data in such a way that valid inferences may be drawn from them, and for extracting the maximum amount of information contained in the data in the most efficient way.

The object of the statistical theory of evolution is to supply a quantitative and mathematical basis for biology in general, and eugenics, the science of man, in particular.

THE THEORY OF SAMPLING DISTRIBUTIONS.

A brief account is given below of Fisher's work on the mathematical theory of statistics to indicate its importance in recent developments in statistics.

The idea of the random sampling distribution of statistics is fundamental in modern statistical theory. The problem of finding such distributions is, however, one of great mathematical difficulty and very little progress was made until Fisher started working on the subject. The earliest example of the modern type of distribution was that of the Chi-square found by Karl Pearson in 1900. Several years later "Student" gave the correct distribution of the sample variance and of his now famous t-statistic or the mean divided by its estimated standard deviation.

I have already mentioned however that Fisher was not acquainted with "Student's" work when he wrote his 1914 paper on the exact distribution of the correlation coefficient. Here he introduced for the first time the brilliant technique of representing a sample of size n by a point in a space of n -dimensions. Such representation has proved extremely useful in subsequent work not only in the theory of distribution, but also in other fields of statistical theory such as the work of J. Neyman and E. S. Pearson. The same representation was used by Fisher in finding the exact distribution of the mean error and the mean square error in 1921,⁷ the regression coefficient in 1922,⁸ the partial correlation coefficient in 1924,⁹ and the coefficient of multiple correlation in 1928.¹⁰

7. *Metron*, i, Part 4, 1-32.

9. *Metron*, iii, 329-332.

8. *Jour. Roy. Stat. Soc.* lxxv, 597-612.

10. *Proc. Roy. Soc. A*, cxxi, 654-673.

In 1923¹¹ he gave a rigorous proof of "Student's" result for the t-statistic, and a little later showed how it could be used for testing various statistical hypotheses.¹² In 1924 Fisher generalized "Student's" t to the well-known z-statistic or half the logarithm of the ratio of two estimated variances based on different degrees of freedom, and was able, with the help of this distribution, to give a unified treatment of practically all the important distributions involved in testing null hypotheses.¹³ The distribution of the discriminant function¹⁴ recently introduced by him to test the significance of the difference, as a whole, of two sets of means of samples drawn from a normal multi-variate correlated population, belongs to the same general class.

The work on the limiting forms of frequency distributions of the largest or smallest member of a sample,¹⁵ of the error of an interpolated value,¹⁶ and the exact distributions which arise in connexion with various tests of significance in harmonic analysis¹⁷ may also be mentioned here.

THEORY OF ESTIMATION AND STATISTICAL INFERENCE.

In science we are always faced with the problem of arguing from the particular to the general, or in statistical language, from the sample to the population. The task of statistical estimation is to find, on the basis of an observed sample, the values of the unknown parameters of the population from which the sample has been derived. Fierce controversy has raged over this subject ever since the publication in 1763 of Bayes' posthumous memoir "An essay towards solving a problem in the doctrine of chances" (*Phil. Trans.* liii, p. 370) in which he proposed to solve this problem with the help of the principle of equal distribution of ignorance. After more than one century and a half new light was shed on this problem by Fisher in his remarkable memoir "On the Mathematical Foundations of Theoretical Statistics."¹⁸ This paper laid the foundations of statistical inference by emphasizing the importance of exact solutions of sampling problems, and by supplying the elements of a valid theory of estimation.

Any statistic¹⁹ which estimates a certain parameter must of course satisfy the criterion of 'consistency', that is, its value must tend to the estimated parameter as the sample size is indefinitely increased. Fisher however gives a second criterion, namely that of 'efficiency', which requires that the variance of the estimating statistic (at least for large samples) should not exceed that of any other consistent statistic estimating the same parameter.

This idea had occurred in an earlier paper of 1920, but a great achievement of the 1921 memoir consisted in providing a machinery for calculating such efficient statistics, in the absence of which we would have been compelled to grope with different consistent statistics, and to compare their variances. This machinery is supplied by Fisher's now well-known method of 'maximum likelihood', a maximum likelihood statistic being always an efficient solution.

In the same memoir Fisher introduces for the first time the idea of a 'sufficient' statistic T , which is inevitably also efficient, and which incorporates the whole of the information available in the sample in regard to a given parameter, in the sense that if T' is any other statistic estimating the same parameter, the joint distribution of T and T' is such that, for a given value of T , the distribution of T' does not involve the parameter estimated. When a sufficient statistic exists, it can always be obtained by the method of maximum likelihood. Many interesting applications of these ideas were given in the same paper, such

11. *Proc. Camb. Phil. Soc.* xci, 655-658.

12. *Econometrica*, iii, 353-365 (1935).

13. *Proc. Camb. Phil. Soc.* xxiv (1928), 180-190.

14. *Proc. Roy. Soc. A*, cxkv (1929), 54-59.

15. *Proc. Int. Math. Congress, Toronto* 1924, 805-813.

16. *Annals of Eugenics*, vii (1936), 179-188.

17. *Proc. Camb. Phil. Soc.* xxiii (1927), 912-921.

18. *Phil. Trans. Roy. Soc. A*, ccxxii, 309-368.

19. The word "statistic" in the singular number has been reserved by Fisher to denote a quantity which is capable of being expressed only in terms of actual observations in the sample. This concept was originally introduced by "Student", and has been given especial emphasis in Fisher's theory.

as errors of grouping, inefficiency of estimation of parameters by the method of moments in Pearson's system of frequency curves, and the estimation of the number of micro-organisms in a sample of water or soil.

The concept of 'information', first tentatively introduced in the 1921 paper, was more fully developed in a later paper²⁰ in 1925, enabling the theory of estimation to be freed from the large sample assumption. The concept of ancillary statistics, which serve to enhance the precision of estimation in the case when no sufficient statistic exists, was also introduced. In 1934 the general method of finding the distribution of any sufficient statistic was indicated,²¹ and it was shown that in certain cases when no sufficient statistic exists, the whole of the information contained in the sample may be recovered by using as ancillary what may be termed the configuration of the sample. The important properties of 'information', as technically defined by Fisher, and 'the maximum likelihood estimate', were clearly summarized in a paper in the same year,²² and a revised proof of the two main theorems in the theory of estimation, together with illustrative examples on the use of ancillary statistics, were given in a paper in 1935.²³

Another important contribution of Fisher to the theory of statistical inference is his idea of 'fiducial probability' introduced in 1930, which is designed to cover the case when we do not want a single estimate of our unknown parameter, but require an interval in which our unknown parameter may be expected to lie.²⁴ Fisher has laid emphasis on the fact that the concept of fiducial probability, though entirely different from that of ordinary probability, is equally rigorous; and has brought out the importance of the fiducial concept in statistical theory in other papers.²⁵

The Chi-square test was perhaps Karl Pearson's greatest single discovery; but Fisher has done much to make its use popular. In 1922 he showed that in the case of contingency tables, for which the margins of the expected table are reconstructed from those of the observed table, the distribution of Chi-square as given by Pearson's formula can be used validly if we take for n' a number exceeding the degrees of freedom by unity.²⁶ This result was discussed in other papers and fully corroborated by material derived from random sampling experiments.²⁷ Fisher also brought the Chi-square test in relation with his general theory of statistical inference, showing in particular that the method of minimising Chi-square agrees for large samples with the method of maximum likelihood.²⁸

There are many other papers in mathematical statistics²⁹ among which the most important are probably those on the use of combinatorial methods for calculating moment coefficients,³⁰ and on the number of reduced Latin Squares of different orders.³¹

STATISTICAL THEORY IN AGRICULTURE.

Fisher's influence in India has been of the greatest importance in connexion with the application of statistical theory to agriculture. The basic principles of the new method are now well-known and need not be discussed in detail. In order to appreciate the revolutionary advance brought about by the introduction of the new technique, let us, however, consider for a moment the contrast between experiments of the old and the new type.

20. *Proc. Camb. Phil. Soc.* xxii, 700-725.

21. *Proc. Roy. Soc. A*, cxliv, 285-307.

22. *Proc. Roy. Soc. A*, cxlvi, 1-8.

23. *Jour. Roy. Stat. Soc.* xcvi, 39-82.

24. *Proc. Camb. Phil. Soc.* xxvi, 528-535.

25. *Proc. Camb. Phil. Soc.* xxviii (1932), 257-261; *Proc. Roy. Soc. A*, cxxxix (1933) 343-348; *Annals Eugenics*, vi (1935), 391-398.

26. *Jour. Roy. Stat. Soc.* lxxv, 87-94.

27. *Economica*, iii (1923), 139-147; *Eugenics Review*, xviii (1926), 32-33.

28. Ref. nos. (18) and (12) and *Bologna. Atti del Congresso Int. dei Matematici*, vi (1928), 94-100.

29. Among other miscellaneous work may be mentioned the method for testing the randomness of a sequence (*Quar. Jour. Met. Soc.* LII, 250, 1926), and the properties and applications of the integrals and derivatives of the normal error function (*Brit. Assoc. Mathematical Tables*, Vol. 1).

30. *Proc. Lond. Math. Soc.* XXX, 199-238 (1929); *Proc. Roy. Soc. A*, cxxx, 16-28. (1930). *Proc. Lond. Math. Soc.* xxxiii, 195-208 (1930).

31. *Proc. Camb. Phil. Soc.* xxx (1934), 492-507.

Suppose we wish to compare the yield of, say, six varieties or the effect on yield of six kinds of manures. In the old type of experiment the field would be divided into six plots, and a single plot would be allotted to each treatment. As Fisher explains, "the treatment giving the highest yield would of course appear to be the best, but no one could say whether the plot would not in fact have yielded as well under some or all of the other treatments."³² It is known that within the same field wide differences exist in the fertility of the soil. Even when the soil fertility is uniform, there are innumerable other causes which affect the yield. How can we be sure that the observed differences in yield are due to the difference in the treatments, and not to soil heterogeneity? How can we be sure that they are not due to chance fluctuations? This is the basic problem. In order to solve it we must eliminate the effect of soil heterogeneity, and make an unbiased estimate of the magnitude of errors due to chance so that we may be sure that the observed effect is significant in comparison with the size of such chance errors.

Let us now see how Fisher solved the problem. Consider the same experimental field which had been originally divided into six portions. Fisher simply further subdivided each portion³³ into a number of plots of smaller size; and within each portion (or block as he called it) he assigned one plot to each treatment but strictly in a random manner. We have now the randomized block in its modern form. Using the principle of block division in two directions symmetrically we get the well-known Latin square.

The important point to be noticed is that the results will be now governed entirely by the laws of chance. There are innumerable causes which produce differences between the plots, and we know from the conditions of the experiment that it is impossible in practice to secure that the plots will be all alike. But the validity of the estimate of error is now guaranteed by the process of randomization, namely "the provision that any two plots, not in the same block, shall have the same probability of being treated alike, and the same probability of being treated differently in each of the ways in which this is possible." The calculus of probability and the apparatus of the statistical theory of sampling distributions can be now used with complete confidence. The logical foundations of scientific inference were thus made secure, and agricultural experiments were placed for the first time on the same footing as experiments in other sciences.

The second point to be observed is that by the technique of block division the problem of soil heterogeneity was solved at the same time. As each block contains all the treatments, differences between the total yields of the different blocks could be safely ascribed, apart from errors of sampling, to soil differences; and could be eliminated by suitable statistical methods. This of course led to a great improvement in the precision of the comparisons. When we remember that in particular experiments in India as much as ninety per cent. of the total variation is sometimes caused by soil differences, the importance of eliminating the effect due to such differences will be easily appreciated.

Replication, randomization and block division or local control are thus the fundamental principles of design introduced by Fisher. Replication is essential because it is the sole source of the estimate of error, while randomization is necessary to guarantee the validity of the estimate, that is, to ensure that the estimate will be unbiased. The purpose of block division is to increase the precision of the comparisons by elimination of soil differences, while replication is also useful in securing the same object by diminishing the experimental error. Finally, the analysis of variance gives a convenient and valid method of extracting the information contained in the observations. As Wishart has pointed out, the Fisherian technique "was something in the nature of a revolution," and altered the subsequent course of agricultural experiments throughout the world.

32. *The Design of Experiments*, 1935, p. 69.

33. I need scarcely add that the experimental field may be divided into any number of convenient portions each of which is further sub-divided into a number of plots.

The modern period of field experiments began in India with the foundation of the Imperial Council of Agricultural Research in 1929, which, from its very inception, laid great emphasis on the use of statistical methods, created a statistical section at its headquarters, and gave a grant to the Statistical Laboratory, Calcutta, for work in this connexion. The earliest analysis of an experiment of the Latin square type was published from the Laboratory in 1931. The first complex (variety-manurial) experiment in 1932, the first split-plot experiment in 1933, and the earliest experiment using the principle of 'confounding' in 1935 were all designed and introduced from the same Laboratory. The use of the new technique has spread rapidly, and no important experiment in India is now laid out on a design of the old type. The ultimate beneficial effect of this movement on Indian agriculture is difficult to estimate or exaggerate.

GENETIC STUDIES.

Fisher's method of maximum likelihood is now well-known and is being increasingly adopted in genetical work in India. More important than these contributions to 'formal questions' as Fisher himself calls it are his contributions to the problems of quantitative characters which depend on a large number of mendelian factors of small individual effects. Almost all economic characters in which the animal or plant breeder is interested are of this nature and here the usual genetic analysis completely breaks down. If, therefore, genetics is to exert its influence on breeding, geneticists and breeders must concentrate on the study of quantitative characters. Fisher has shown how statistics of the second and third degree can be utilised for this purpose, and has given methods for the estimation of various genetic quantities from which genetic variances can be calculated and the effects of dominance and environmental causes eliminated.

A knowledge of genetic variance and of the possible number of factors involved in a character is of fundamental importance to the breeder in assessing the selection potentiality of his material. Fisher has shown the way to attain this knowledge. The experimental study, however, requires a sound lay out and elaborate recording of observational data to make statistical analysis possible. Work has recently been started on cotton at the Institute of Plant Industry at Indore on the new lines, and it may be confidently asserted that the influence of Fisher will ere long become increasingly important in this field of study also.

THE INFLUENCE OF FISHER ON STATISTICAL WORK IN INDIA.

In recent developments in statistics in India the most powerful influence has been that of R. A. Fisher. The movement for the study of analytic statistics started in this country at a time when Fisher's work was just coming into prominence, and the younger workers have been all brought up in what may be called the Fisherian tradition. The methods developed by him are being used not only in agriculture, but also in all kinds of investigations of importance to national welfare. On the theoretical side, what little is being done in India is based largely on his work. He has been the accepted and acknowledged leader for a long time, but his presence here as the first President of the Indian Statistical Conference has established personal relations which are highly cherished by all statistical workers in India.*

P. C. MAHALANOBIS.

CALCUTTA,
6th January, 1938.

* Written on the occasion of the opening of the First Session of the Indian Statistical Conference presided over by Prof. R. A. Fisher in Calcutta on the 7th January, 1938.