

A NOTE ON A TRANSPLANTATION THEOREM OF KANJIN AND MULTIPLE LAGUERRE EXPANSIONS

S. THANGAVELU

(Communicated by J. Marshall Ash)

ABSTRACT. By applying a transplantation theorem of Kanjin, a multiplier theorem and a Cesàro summability result are proved for multiple Laguerre expansions. In the one-dimensional case an improved version of the multiplier theorem is obtained.

1

Consider the normalised Laguerre functions \mathcal{L}_k^α , $\alpha > -1$, on $\mathbb{R}_+ = (0, \infty)$ defined by

$$(1.1) \quad \mathcal{L}_k^\alpha(t) = \left(\frac{\Gamma(k+1)}{\Gamma(k+\alpha+1)} \right)^{1/2} L_k^\alpha(t) e^{-t/2} t^{\alpha/2}$$

where $L_k^\alpha(t)$ are the Laguerre polynomials of type α . The functions $\{\mathcal{L}_k^\alpha\}$ form a complete orthonormal system for $L^2(\mathbb{R}_+)$. Recently, in [4] Kanjin studied the mapping properties of the operator T_α^β , which is defined as

$$(1.2) \quad T_\alpha^\beta f = \sum_{k=0}^{\infty} (f, \mathcal{L}_k^\beta) \mathcal{L}_k^\alpha$$

where (f, g) stands for the inner product in $L^2(\mathbb{R}_+)$. For the operator T_α^β he has proved the following result.

Theorem 1.1 (Kanjin). *Let $\alpha, \beta > -1$ and $\gamma = \min\{\alpha, \beta\}$. If $\gamma \geq 0$ then*

$$(1.3) \quad \|T_\alpha^\beta f\|_p \leq C \|f\|_p \quad \text{for } 1 < p < \infty.$$

If $-1 < \gamma < 0$ then (1.3) is valid for p in the interval $(1 + \gamma/2)^{-1} < p < -2/\gamma$.

The above theorem is called a transplantation theorem for the following reason. Given a bounded sequence $\lambda(k)$ we can define an operator M_λ^α by setting

$$(1.4) \quad M_\lambda^\alpha f = \sum_{k=0}^{\infty} \lambda(k) (f, \mathcal{L}_k^\alpha) \mathcal{L}_k^\alpha$$

whenever f has the Laguerre expansion

$$(1.5) \quad f = \sum_{k=0}^{\infty} (f, \mathcal{L}_k^\alpha) \mathcal{L}_k^\alpha.$$

From the theorem, we can deduce the norm inequality

$$(1.6) \quad \|M_\lambda^\alpha f\|_p \leq C \|f\|_p$$

for any α if we know (1.6) for a particular α_0 . This follows from the identity

$$(1.7) \quad T_\beta^\alpha M_\lambda^\alpha T_\alpha^\beta f = M_\lambda^\beta f.$$

As an application Kanjin proves the following result concerning M_λ^α .

Theorem 1.2 (Kanjin). *Let $\lambda(t)$ be a four times differentiable function on $(0, \infty)$ and satisfy*

$$(1.8) \quad \sup_{t>0} |t^k \lambda^{(k)}(t)| \leq c_k$$

for $k = 0, 1, 2, 3, 4$. Then (1.6) is true for $1 < p < \infty$ if $\alpha \geq 0$ and for $(1 + \alpha/2)^{-1} < p < -2/\alpha$ if $-1 < \alpha < 0$.

Theorem 1.2 is deduced by applying the transplantation theorem to the particular case $\alpha = 0$, which is proved by Dlugosz in [1]. Now the aim of this note is to prove an improved version of the above multiplier theorem and also to give applications to higher-dimensional Laguerre expansions.

2

Let $\mathbb{R}_+^n = \{x \in \mathbb{R}^n : x_j \geq 0 \text{ for all } j\}$, and consider for every $\alpha \in \mathbb{R}_+^n$ and a multi-index $m = (m_1, m_2, \dots, m_n)$, the normalised Laguerre functions \mathcal{L}_m^α on \mathbb{R}_+^n defined by

$$(2.1) \quad \mathcal{L}_m^\alpha(x) = \prod_{j=1}^n \mathcal{L}_{m_j}^{\alpha_j}(x_j).$$

They form a complete orthonormal system for $L^2(\mathbb{R}_+^n)$, and the Laguerre expansion of a function f in $L^p(\mathbb{R}_+^n)$ can be written as

$$(2.2) \quad f = \sum_{m=0}^{\infty} (f, \mathcal{L}_m^\alpha) \mathcal{L}_m^\alpha$$

where the sum is extended over all the multi-indices. Expansions of the above type have been studied by Dlugosz [1] when α is a multi-index.

For the above series (2.2) we define the Cesàro means σ_N^δ of order δ by the equation

$$(2.3) \quad \sigma_N^\delta f = \frac{1}{A_N^\delta} \sum_{k=0}^N A_{N-k}^\delta \sum_{|m|=k} (f, \mathcal{L}_m^\alpha) \mathcal{L}_m^\alpha$$

where $A_k^\delta = \Gamma(k+\delta+1)/\Gamma(k+1)$ are the binomial coefficients. Given a function λ on $(0, \infty)$ we also define the multiplier operator M_λ^α as

$$(2.4) \quad M_\lambda^\alpha f = \sum_{k=0}^{\infty} \lambda(2k+n) \sum_{|m|=k} (f, \mathcal{L}_m^\alpha) \mathcal{L}_m^\alpha.$$

For the operators (2.3) and (2.4) we prove the following two theorems.

Theorem 2.1. Let $\delta > \frac{1}{2}$. Then the uniform estimates

$$(2.5) \quad \|\sigma_N^\delta f\|_p \leq C\|f\|_p$$

are valid iff $4n/(2n+1+2\delta) < p < 4n/(2n-1-2\delta)$.

Theorem 2.2. Assume that the function λ satisfies the conditions

$$(2.6) \quad \sup_{t>0} |t^k \lambda^{(k)}(t)| \leq c_k$$

for $k = 0, 1, 2, \dots, \nu$ where $\nu = n+1$ if n is odd and $\nu = n+2$ if n is even. Then for $1 < p < \infty$ we have

$$(2.7) \quad \|M_j^\alpha f\|_p \leq C\|f\|_p.$$

In the case $n = 1$ we can take $\nu = 1$ in the hypothesis and (2.7) is valid for $\frac{4}{3} < p < 4$.

A slightly weaker form of Theorem 2.2 is proved in [1] when α is a multi-index. In that version one has $\nu = n+3$ for all n . Theorem 2.1 is known when $n = 1$ and is due to Gorlich and Markett [3, 5].

For the Laguerre series (2.2) we also define the Riesz transforms R_j , $j = 1, 2, \dots, n$, by the formula

$$(2.8) \quad R_j f = \sum_{m=0}^{\infty} (2m_j + 1)(2|m| + n)^{-1} (f, \mathcal{L}_m^\alpha) \mathcal{L}_m^\alpha.$$

Riesz transforms for the Hermite and special Hermite expansions have been studied by the author in [9, 12]. For the above Riesz transforms (2.8) we prove

Theorem 2.3. For $1 < p < \infty$ all the Riesz transforms R_j are bounded on $L^p(\mathbb{R}_+^n)$.

All three theorems will be proved by appealing to the n -dimensional version of Kanjin's transplantation Theorem 1.1. For α, β in \mathbb{R}_+^n we define T_α^β by

$$(2.9) \quad T_\alpha^\beta f = \sum_{m=0}^{\infty} (f, \mathcal{L}_m^\beta) \mathcal{L}_m^\alpha.$$

Then, for f in $C_0^\infty(\mathbb{R}_+^n)$ and $1 < p < \infty$,

$$(2.10) \quad \|T_\alpha^\beta f\|_p \leq C\|f\|_p.$$

This follows from Theorem 1.1 by iteration.

In view of (2.10) Theorems 2.1, 2.2, and 2.3 will follow once we show that they are true in the particular case $\alpha = 0$. It will be shown in the next section that the case $\alpha = 0$ follows from known results on special Hermite expansions as a special case. The one-dimensional case of Theorem 2.2 when $\alpha = \frac{1}{2}$ will be deduced from the corresponding result on the Hermite expansions. This will be done in the last section.

3

Consider the functions $\psi_m(z)$ on \mathbb{C}^n defined by

$$(3.1) \quad \psi_m(z) = \prod_{j=1}^n L_{m_j}(\frac{1}{2}|z_j|^2) e^{-|z_j|^2/4}$$

where $L_k(t)$ are the Laguerre polynomials of type 0. The functions $\psi_m(z)$ are called special Hermite functions since they are related to the Hermite function $\Phi_m(x)$ on \mathbb{R}^n . This terminology is due to Strichartz [6]. In fact, one has

$$(3.2) \quad \psi_m(z) = \int_{\mathbb{R}^n} e^{i\lambda \cdot \xi} \Phi_m\left(\xi + \frac{y}{2}\right) \Phi_m\left(\xi - \frac{y}{2}\right) d\xi$$

where $z = x + iy$, $x, y \in \mathbb{R}^n$ (see [2]). Given f on \mathbb{C}^n we have the special Hermite expansion

$$(3.3) \quad f(z) = (2\pi)^{-n} \sum_{m=0}^{\infty} f \times \psi_m(z)$$

where the twisted convolution $f \times g$ of two functions is defined by

$$(3.4) \quad f \times g(z) = \int_{\mathbb{C}^n} f(z-w)g(w)e^{(i/2)\text{Im } z \cdot \bar{w}} dw.$$

We can also write (3.3) in the form

$$(3.5) \quad f(z) = (2\pi)^{-n} \sum_{k=0}^{\infty} f \times \varphi_k^{n-1}(z)$$

where $\varphi_k^{n-1}(z) = L_k^{n-1}(\frac{1}{2}|z|^2)e^{-|z|^2/4}$. For all these facts we refer to [11].

For the special Hermite expansion let C_N^δ be the Cesàro means defined by

$$(3.6) \quad C_N^\delta f = \frac{1}{A_N^\delta} \sum_{k=0}^N A_{N-k}^\delta \sum_{|m|=k} (f \times \psi_m).$$

Given a function λ on $(0, \infty)$ we also define a multiplier transform T_λ by

$$(3.7) \quad T_\lambda f = \sum_{k=0}^{\infty} \lambda(2k+n) \sum_{|m|=k} (f \times \psi_m).$$

In [11] we proved

Theorem 3.1. *Let $\delta > \frac{1}{2}$. Then for f in $L^p(\mathbb{C}^n)$*

$$\|C_N^\delta f\|_p \leq C \|f\|_p$$

holds if and only if $4n/(2n+1+2\delta) < p < 4n/(2n-1-2\delta)$.

Regarding T_λ we have proved the following multiplier theorem in [10].

Theorem 3.2. *Let λ satisfy the hypothesis of Theorem 2.2. Then for $1 < p < \infty$ one has $\|T_\lambda f\|_p \leq C \|f\|_p$.*

The case $\alpha = 0$ of Theorems 2.1 and 2.2 will be deduced from the above theorems in the following way. When f is a radial function the twisted convolution $f \times \varphi_k^{n-1}$ becomes

$$(3.8) \quad f \times \varphi_k^{n-1}(z) = \frac{k!(n-1)!}{(k+n-1)!} \left(\int_0^\infty f(r) \varphi_k^{n-1}(r) r^{2n-1} dr \right) \varphi_k^{n-1}(z)$$

where $\varphi_k^{\alpha-1}(r) = \varphi_k^{\alpha-1}(z)$ with $|z| = r$. If f is a polyradial function, i.e., $f(z_1, \dots, z_n) = f(r_1, \dots, r_n)$, $r_j = |z_j|$, then in view of (3.8) and (3.1) one has

$$(3.9) \quad f \times \psi_m = \left\{ \int_{\mathbb{R}_+^n} f(r_1, \dots, r_n) \left(\prod_{j=1}^n \mathcal{L}_m(\frac{1}{2}r_j^2) \right) r_1 \cdots r_n dr_1 \cdots dr_n \right\} \psi_m.$$

Therefore, one sees that

$$(3.10) \quad f \times \psi_m(\sqrt{2}z) = (g, \mathcal{L}_m)\mathcal{L}_m(r)$$

where $g(r_1, \dots, r_n) = f(\sqrt{2}r_1, \dots, \sqrt{2}r_n)$. Therefore, $C_N^\delta f$ becomes $\sigma_N^\delta g$ and $T_\lambda f$ becomes $M_\lambda^\delta g$; hence, the case $\alpha = 0$ of Theorems 2.1 and 2.2 follow.

The case $\alpha = 0$ of Theorem 3.3 follows from the fact (see [12]) that the Riesz transforms

$$(3.11) \quad S_j f = \sum_{m=0}^\infty (2m_j + 1)(2|m| + n)^{-1} f \times \psi_m$$

for the special Hermite expansions are bounded on $L^p(\mathbb{C}^n)$, $1 < p < \infty$.

4

Consider the normalised Hermite functions $h_k(x)$ on \mathbb{R} . We also consider the Laguerre function φ_k^α of another type defined by, for α real,

$$(4.1) \quad \varphi_k^\alpha(x) = \mathcal{L}_k^\alpha(x^2)(2x)^{1/2}, \quad x \in \mathbb{R}_+.$$

Then the Hermite functions h_k and φ_k^α are related by (see [7])

$$(4.2) \quad h_{2k}(x) = (-1)^k \frac{1}{\sqrt{2}} \varphi_k^{-1/2}(x), \quad h_{2k+1}(x) = (-1)^k \frac{1}{\sqrt{2}} \varphi_k^{1/2}(x).$$

Consider a multiplier transform M for the Hermite series defined by

$$(4.3) \quad Mf(x) = \sum_{k=0}^\infty \lambda(k)(f, h_k)h_k(x).$$

In [8] we proved

Theorem 4.1. *Assume that λ is bounded and satisfies $|t\lambda'(t)| \leq C$ for all $t > 0$. Then M is bounded on $L^p(\mathbb{R})$, $1 < p < \infty$.*

Since h_{2k} is even and h_{2k+1} is odd, by considering f to be odd we see that

$$(4.4) \quad Mf(x) = \sum_{k=0}^\infty \lambda(2k+1)(f, \varphi_k^{1/2})\varphi_k^{1/2}(x),$$

and this is related to $M_\lambda^{1/2}$ in the following way. An easy calculation shows that

$$(4.5) \quad (f, \varphi_k^{1/2}) = \frac{1}{\sqrt{2}}(g, \mathcal{L}_k^{1/2})$$

where $f(\sqrt{x})x^{-1/4} = g(x)$. Therefore,

$$(4.6) \quad Mf(\sqrt{x})x^{-1/4} = 2 \sum_{k=0}^{\infty} \lambda(2k+1)(g, \mathcal{L}_k^{1/2}) \mathcal{L}_k^{1/2}(x).$$

If we know that for $\frac{4}{3} < p < 4$

$$(4.7) \quad \int_0^{\infty} |Mf(x)|^p x^{-p/2+1} dx \leq C \int_0^{\infty} |f(x)|^p x^{-p/2+1} dx$$

then it follows that

$$(4.8) \quad \int_0^{\infty} |M_{\lambda}^{1/2} g(x)|^p dx \leq C \int_0^{\infty} |g(x)|^p dx;$$

hence, the case $n = 1$, $\alpha = \frac{1}{2}$ of Theorem 2.2 follows. We claim that (4.7) is true.

To prove the claim we recall the proof of Theorem 4.1. Let T^t be the semigroup on $L^p(\mathbb{R})$ defined by

$$(4.9) \quad T^t f = \sum_{k=0}^{\infty} e^{-(2k+1)t} (f, h_k) h_k.$$

For this semigroup we defined the g and g^* functions in the following way:

$$(4.10) \quad (g(f, x))^2 = \int_0^{\infty} t |\partial_t T^t f(x)|^2 dt,$$

$$(4.11) \quad (g^*(f, x))^2 = \int_{-\infty}^{\infty} \int_0^{\infty} t^{1/2} (1 + t^{-1/2}|x-y|)^{-2} |\partial_t T^t f(y)|^2 dy dt.$$

For the g and g^* functions we proved that

$$(4.12) \quad C_1 \|f\|_p \leq \|g(f)\|_p \leq C_2 \|f\|_p, \quad 1 < p < \infty,$$

$$(4.13) \quad \|g^*(f)\|_p \leq C \|f\|_p, \quad p > 2.$$

Under the assumption that $|t\lambda'(t)|$ is bounded we verified that

$$(4.14) \quad g(Mf, x) \leq C g^*(f, x),$$

and in view of (4.12) and (4.13) this proved Theorem 4.1.

Therefore, in order to prove the weighted version we need to check that

$$(4.12)' \quad C_1 \|f\|_{p,w} \leq \|g(f)\|_{p,w} \leq C_2 \|f\|_{p,w}, \quad \frac{4}{3} < p < 4,$$

$$(4.13)' \quad \|g^*(f)\|_{p,w} \leq C \|f\|_{p,w}, \quad 2 < p < 4,$$

where $\|f\|_{p,w}$ stands for the norm

$$\|f\|_{p,w} = \left(\int_{-\infty}^{\infty} |f(x)|^p |x|^{-p/2+1} dx \right)^{1/p}.$$

Thus we need weighted norm inequalities for the g and g^* functions.

In [8] we proved the L^p boundedness of g by applying singular integral theory. We identified g with a singular integral operator whose kernel takes

values in the Hilbert space $L^2(\mathbb{R}_+, t dt)$. When the weight function w is in the Muckenhoupt class A_p (see [13]) then we also have

$$(4.15) \quad \int_{-\infty}^{\infty} |g(f)|^p w(x) dx \leq C \int_{-\infty}^{\infty} |f(x)|^p w(x) dx.$$

When $\frac{4}{3} < p < 4$, $w(x) = |x|^{-p/2+1}$ is in A_p ; hence, the right-hand side inequality of (4.12)' is valid. We will now show that the reverse inequality is also valid.

From [8] we recall that we have the partial isometry

$$(4.16) \quad \|g(f)\|_2 = \frac{1}{2} \|f\|_2;$$

from this, by polarisation, we obtain

$$(4.17) \quad \left| \int_{-\infty}^{\infty} f_1(x) \bar{f}_2(x) dx \right| = 4 \int_{-\infty}^{\infty} \int_0^{\infty} t \partial_t T^t f_1(x) \overline{\partial_t T^t f_2(x)} dt dx.$$

This gives the inequality

$$(4.18) \quad \left| \int_{-\infty}^{\infty} f_1(x) \bar{f}_2(x) dx \right| \leq 4 \int_{-\infty}^{\infty} g(f_1, x) g(f_2, x) dx.$$

Let us now take $h(x) = f_2(x)|x|^{-1/2+1/p}$ so that

$$(4.19) \quad \left| \int_{-\infty}^{\infty} f_1(x) |x|^{-1/2+1/p} \bar{f}_2(x) dx \right| \leq 4 \int_{-\infty}^{\infty} g(f_1, x) |x|^{-1/2+1/p} g(h, x) |x|^{-1/2+1/q} dx$$

where q is the index conjugate to p . An application of Holder's inequality gives

$$(4.20) \quad \int_{-\infty}^{\infty} g(f_1, x) |x|^{-1/2+1/p} g(h, x) |x|^{-1/2+1/q} dx \leq \left(\int_{-\infty}^{\infty} |g(f_1, x)|^p |x|^{-p/2+1} dx \right)^{1/p} \left(\int_{-\infty}^{\infty} |g(h, x)|^q |x|^{-q/2+1} dx \right)^{1/q}.$$

Applying the direct inequality (4.12)' to the second factor we get

$$(4.21) \quad \int_{-\infty}^{\infty} |g(h, x)|^q |x|^{-q/2+1} dx \leq C \int_{-\infty}^{\infty} |f_2(x)|^q |x|^{-q/2+q/p+1-q/2} dx \leq C \int_{-\infty}^{\infty} |f_2(x)|^q dx.$$

In view of (4.20) and (4.21) the inequality (4.19) becomes

$$(4.22) \quad \left| \int_{-\infty}^{\infty} f_1(x) |x|^{-1/2+1/p} \bar{f}_2(x) dx \right| \leq C \|g(f_1)\|_{p,w} \|f_2\|_q.$$

Taking the supremum over all f with $\|f_2\|_q \leq 1$ we obtain

$$(4.23) \quad \int_{-\infty}^{\infty} |f_1(x)|^p |x|^{-p/2+1} dx \leq C \|g(f_1)\|_{p,w}.$$

This completes the proof of (4.12)'.

To establish the inequality (4.13)' we observe that

$$(4.24) \quad \int_{-\infty}^{\infty} (g^*(f, x))^2 h(x) dx \leq \int_{-\infty}^{\infty} (g(f, x))^2 \Lambda h(x) dx$$

for every nonnegative function h where Λh is the Hardy-Littlewood maximal function. If $2 < p < 4$, let $r = p/2$ and s be the conjugate index of r . Setting $h_1(x) = h(x)|x|^{-1+1/r}$ we have

$$(4.25) \quad \begin{aligned} & \int_{-\infty}^{\infty} (g^*(f, x))^2 |x|^{-1+1/r} h(x) dx \\ & \leq C \int_{-\infty}^{\infty} (g(f, x))^2 |x|^{-1+1/r} |x|^{1/s} \Lambda h_1(x) dx \\ & \leq C \left(\int_{-\infty}^{\infty} (g(f, x))^p |x|^{-p/2+1} dx \right)^{2/p} \left(\int_{-\infty}^{\infty} |x| (\Lambda h_1(x))^s ds \right)^{1/s} \end{aligned}$$

by an application of Holder's inequality. Since $s > 2$, $|x| \in A_s$; hence,

$$(4.26) \quad \begin{aligned} \int_{-\infty}^{\infty} |x| (\Lambda h_1(x))^s ds & \leq C \int_{-\infty}^{\infty} |h(x)|^s |x|^{-s+s/r+1} dx \\ & \leq C \int_{-\infty}^{\infty} |h(x)|^s dx. \end{aligned}$$

Thus we have the inequality

$$(4.27) \quad \begin{aligned} & \int_{-\infty}^{\infty} (g^*(f, x))^2 |x|^{-1+1/r} h(x) dx \\ & \leq C \left(\int_{-\infty}^{\infty} |f(x)|^p |x|^{-p/2+1} dx \right)^{2/p} \|h\|_s. \end{aligned}$$

Taking the supremum over all h with $\|h\|_s \leq 1$ we obtain

$$(4.28) \quad \int_{-\infty}^{\infty} (g^*(f, x))^p |x|^{-p/2+1} dx \leq C \int_{-\infty}^{\infty} |f(x)|^p |x|^{-p/2+1} dx.$$

This proves the inequality (4.13)'.

Therefore, in view of (4.12)', (4.13)', and (4.14) we obtain the weighted inequality

$$(4.29) \quad \int_{-\infty}^{\infty} |Mf(x)|^p |x|^{-p/2-1} dx \leq C \int_{-\infty}^{\infty} |f(x)|^p |x|^{-p/2+1} dx$$

for $\frac{4}{3} < p < 4$, and this proves the multiplier theorem for $\alpha = \frac{1}{2}$. By applying the transplantation theorem we complete the proof of Theorem 2.2 when $n = 1$.

REFERENCES

1. D. Dlugosz, L^p multipliers for the Laguerre expansions, *Colloq. Math.* **54** (1987), 287-293.
2. G. Folland, *Harmonic analysis in phase space*, *Ann. of Math. Stud.*, vol. 112, Princeton Univ. Press, Princeton, NJ, 1989.
3. E. Gorlich and C. Markett, *Mean Cesaro summability and operator norms for Laguerre expansions*, *Comment. Math. Prace Mat. Tomus Specialus II* (1979), 139-148.
4. Y. Kanjin, *A transplantation theorem for Laguerre series*, preprint, 1990.

5. C. Markett, *Norm estimates for Cesaro means of Laguerre expansions*, Approximation and Function Spaces (Proc. Conf. Gdansk, 1979), North-Holland, Amsterdam, 1981, pp. 419–435.
6. R. Strichartz, *Harmonic analysis as spectral theory of Laplacians*, J. Funct. Anal. **87** (1989), 51–148.
7. G. Szego, *Orthogonal polynomials*, Amer. Math. Soc. Colloq. Publ., vol. 23, Amer. Math. Soc., Providence, RI, 1967.
8. S. Thangavelu, *Multipliers for Hermite expansions*, Mat. Ibero Americana **3** (1987), 1–24.
9. —, *Riesz transforms and the wave equation for the Hermite operator*, Comm. Partial Differential Equations **15** (1990), 1199–1215.
10. —, *Littlewood-Paley-Stein theory on \mathbb{C}^n and Weyl multipliers*, Rev. Mat. Ibero Americana **6** (1990), 75–90.
11. —, *Bochner-Riesz means, Weyl multipliers and special Hermite expansions*, Ark. Mat. **29** (1991), 307–321.
12. —, *Conjugate Poisson integrals and Riesz transforms for Hermite and special Hermite expansions*, Colloq. Math. **64** (1993), 103–113.
13. A. Torchinsky, *Real variable methods in harmonic analysis*, Academic Press, London, 1986.